

Ongoing Cardiovascular Outcomes Trial (CVOT) Timeline - July 19, 2018

The table below includes an overview of ongoing cardiovascular outcomes trials (CVOTs). It includes
all of the ongoing CVOTs related to diabetes drugs or PCSK9 inhibitors that we are aware of, though
we acknowledge that it may be incomplete. We will continuously update the table as timelines
change.

Molecule	Name of Clinical Trial	Class of Drug	Study Enrollment	Primary Outcome Measures	Expected Completion Date
Tradjenta/Trajenta (linagliptin) vs. glimepiride	CAROLINA	DPP-4 inhibitor vs. sulfonylurea	6,115	4-point MACE composite endpoint (CV death, nonfatal MI, nonfatal stroke, and hospitalization for unstable angina)	August 2018
Steglatro (ertugliflozin)	VERTIS CV	SGLT-2 inhibitor	8,000	3-point MACE composite endpoint (CV death, non- fatal MI, non- fatal stroke)	September 2019
Farxiga (dapagliflozin) in HFrEF	Dapa-HF	SGLT-2 inhibitor	4,744	composite of CV death, hospitalization for heart failure, or urgent HF visit	December 2019
Jardiance (empagliflozin) in HFpEF	EMPEROR- Preserved	SGLT-2 inhibitor	4,126	composite of CV death or hospitalization for heart failure	June 2020

www.closeconcerns.com 1

Jardiance (empagliflozin) in HFrEF	EMPEROR- Reduced	SGLT-2 inhibitor	2,850	composite of CV death or hospitalization for heart failure	June 2020
Farxiga (dapagliflozin) in CKD	Dapa-CKD	SGLT-2 inhibitor	4,000	Composite of (i) ≥50% sustained decline in eGFR, (ii) reaching ESRD, and (iii) CV or renal death	November 2020
sotagliflozin	SOLOIST- WHF	SGLT-1/2 dual inhibitor	4,000	composite of CV death or hospitalization for heart failure	January 2021
Farxiga (dapagliflozin) in HFpEF	DELIVER	SGLT-2 inhibitor	4,700	composite of CV death, hospitalization for heart failure, or urgent HF visit	June 2021
sotagliflozin	SCORED	SGLT-1/2 dual inhibitor	10,500	3-point MACE composite (CV death, non- fatal MI, non- fatal stroke); CV death/ hospitalization for heart failure composite	March 2022
Jardiance (empagliflozin) in CKD	EMPA- KIDNEY	SGLT-2 inhibitor	5,000	Composite of (i) CKD progression - ESKD, decline in eGFR to <10 mL/min/ 1.73m², renal death, or a	June 2022

www.closeconcerns.com 2

				sustained decline of ≥40% in eGFR from randomization, and (ii) CV death	
semaglutide for obesity	SELECT	GLP-1 agonist	17,500	3-point MACE composite (CV death, non- fatal MI, non- fatal stroke)	September 2023
oral semaglutide	"SOUL 2"	oral GLP-1 agonist	unknown	unknown	unknown

www.closeconcerns.com 3